

NORMAL GENERATION OF LOCALLY COMPACT GROUPS

A. EISENMANN AND N. MONOD

ABSTRACT. It is a well-known open problem since the 1970s whether a finitely generated perfect group can be normally generated by a single element or not. We prove that the topological version of this problem has an affirmative answer as long as we exclude infinite discrete quotients (which is probably a necessary restriction).

1. INTRODUCTION

Let G be a perfect group. Can G be *normally generated* by a single element?

Equivalently: can G be made trivial by adding a single relation? In other words yet: does G contain some element not belonging to any proper normal subgroup?

This is well-known to be true when G is *finite* (see e.g. [4, 4.2] for a stronger fact). For infinite groups, it is a long-standing open problem attributed to J. Wiegold; see [1, FP14] and [5, 5.52]. The group G is assumed to be finitely generated in order to avoid obvious counter-examples (infinite direct sums of perfect groups). It is widely believed to be false: for instance, Lennox and Wiegold conjecture that a free product of n finite groups cannot be normally generated by less than $n/2$ elements [4]. A completely different potential source of counter-examples is proposed in [6].

The question can be asked more generally for topological groups G (which we always suppose Hausdorff). Here G is called *perfect* if it has no non-trivial continuous homomorphism to an abelian topological group, or equivalently if its commutator subgroup is dense. Likewise, it is deemed *normally generated* by a single element $g \in G$ if the smallest closed normal subgroup containing g is G itself.

If G has an infinite discrete quotient group, then the putative counter-examples for abstract groups will of course be *a fortiori* counter-examples for G itself. Rather surprisingly, the question has a positive answer when we exclude such quotients:

Theorem 1. *Let G be a compactly generated locally compact group without infinite discrete quotient. If G is perfect, then it is normally generated by a single element.*

This result uses structure theory, both classical [3, 7] and very recent [2].

Remark 2. Compact generation is the replacement for the necessary assumption of finite generation. It remains necessary even when there is no infinite discrete quotient, as shown by an example constructed for us by Y. de Cornulier, see Section 3 below.

Acknowledgements. Our motivation for this work came from discussions with Taka Ozawa and Andreas Thom, as always pleasant and inspiring. We are grateful to Yves de Cornulier for allowing us to present his example for Remark 2.

Supported in part by the European Research Council and the Swiss National Science Foundation.

2. PROOF OF THE THEOREM

We begin with a few simple facts. It is rather straightforward to go from finite groups to profinite groups:

Lemma 3. *A perfect profinite group is normally generated by a single element.*

Proof. Let $G = \varprojlim G/G_\alpha$, be a perfect profinite group, where G_α runs over the open normal subgroups of G . Each G/G_α is a perfect finite group and thus it is normally generated by a single element. Denote by X_α the set of elements of G whose image normally generates G/G_α . Thus X_α is a non-empty and clopen subset of G . The family of these sets is directed and thus its intersection is non-empty by compactness of G . We claim that any element g in this intersection normally generates G . Indeed, suppose that g is contained in a closed normal subgroup N of G . In a profinite group, every closed subgroup is an intersection of open subgroups. Since moreover open subgroups have finitely many conjugates, we see that N is an intersection of open normal subgroups. By construction, g does not belong to any $G_\alpha \neq G$. It follows $N = G$, as claimed. \square

We shall repeatedly but tacitly use the following fact: if S is a subset of a topological group G and $N \triangleleft G$ is a closed normal subgroup, then the image of S in G/N is dense (if and) only if SN is dense in G . Indeed, the quotient map $G \rightarrow G/N$ is open.

Lemma 4. *Let G be a perfect topological group, and let N be a closed normal soluble subgroup of G . If G/N is normally generated by a single element, then so is G .*

Proof. Let $g \in G$ be such that its image in the quotient G/N normally generates it. Let H be the minimal closed normal subgroup of G containing g . Then HN is dense in G . Thus, the canonical homomorphism $N \rightarrow G/H$ has dense image, which implies that G/H is soluble. Being also perfect, it must be trivial. Thus $H = G$ as needed. \square

We endow the product of any family of topological groups with the product topology. We recall that all but finitely many factors must be compact if the product is locally compact — for instance, if it appears as a closed subgroup of a locally compact group. In order to use structure theory, we need the following.

Proposition 5. *Let G be a perfect topological group. Let H be a closed normal subgroup of G which is the topological direct product of topologically simple topological groups. If G/H is generated by a single element, then so is G .*

Proof. Let $H = \prod_{i \in I} S_i$, where S_i is a simple topological group for each i in some index set I . We can assume that no S_i is abelian by applying Lemma 4 to the product of all abelian factors S_i . Let N be any closed normal subgroup of G . We denote by $[N, H]$ the closed subgroup generated by commutators $[n, h]$ with $n \in N$ and $h \in H$. Hence $[N, H]$ is a closed normal subgroup of G and thus also of H . Therefore, there is a subset J of I such that $[N, H] = \prod_{i \in J} S_i$. Let $H_1 := \prod_{i \in I \setminus J} S_i$. Then $H = [N, H] \times H_1$ and the action of N by conjugation induces a permutation on I , permuting the simple subgroups S_i . As $[N, H]$ is normalized by N , this permutation action on I splits into actions on J and on $I \setminus J$. In particular, N normalizes H_1 . Thus $[N, H_1] \subset H_1$. But on the other hand $[N, H_1] \subset [N, H]$ and thus $[N, H_1]$ is trivial since $[N, H] \cap H_1$ is trivial. In other words, N centralizes H_1 .

Let now g be an element of G whose image in G/H is normally generating. We apply the above argument to the smallest closed normal subgroup N of G containing g . This yields a

dense normal subgroup $NH = NH_1$ in G , with N and H_1 centralizing each other. Choose an element $h \in H_1$ that is non-trivial in each coordinate S_i when $i \in I \setminus J$. Thus the group H_1 is normally generated by h by a standard argument: if $L \triangleleft H_1$ is the minimal closed normal subgroup containing h , then for all $i \in I \setminus J$, $[L, S_i]$ is a non-trivial closed normal subgroup of S_i and hence equals S_i . Therefore L contains S_i and thus indeed $L = H_1$.

We claim that gh normally generates G . Let thus $M \triangleleft G$ be the minimal closed normal subgroup containing gh . The image of M in G/N contains $ghN = hN$. Since the image of H_1 in G/N is dense, we conclude that the image of M is dense too. That is, NM is dense in G . Turning things around, the image of N in G/M is dense and therefore the image of H_1 in G/M is central. Since H_1 is perfect, this image of H_1 is in fact trivial, and thus $h \in M$. It now follows $g \in M$, hence $N < M$ and thus $NM = M$ which finally implies $M = G$ as claimed. \square

We now prove Theorem 1. Let G be a compactly generated locally compact group without infinite discrete quotients. We begin with some classical structure theory. Because of the solution of Hilbert's fifth problem, there exists a maximal normal compact subgroup K in the identity component G° of G and the quotient G°/K is a connected Lie group; see Lemma 2.2 in [2]. Since K is necessarily unique, it is also characteristic in G . Accordingly, its identity component K° , the commutator subgroup $[K^\circ, K^\circ]$ and the centre $Z([K^\circ, K^\circ])$ are all characteristic in G . Therefore, in view of Lemma 4, we can assume that $Z([K^\circ, K^\circ])$ is trivial. In that case, $[K^\circ, K^\circ]$ is a product of (possibly infinitely many) simple compact Lie groups; see Theorem 9.19 in [3] (recalling that the commutator of a compact connected group is "semisimple" in the terminology of [3]). Since these simple Lie groups are centre-free, they are topologically simple. In view of Proposition 5, we can therefore assume that $[K^\circ, K^\circ]$ itself is trivial. Now K° is abelian and can be assumed trivial by Lemma 4. We deduce that K is central in G° since any totally disconnected normal subgroup of a connected group is central; appealing one more time to Lemma 4, we can assume K trivial. In other words, G° is a Lie group. Applying Lemma 4, we can assume that its radical as well as its centre are trivial. Therefore G° is a centre-free semi-simple connected Lie group, and thus a product of topologically simple groups. Proposition 5 shows that we can assume it to be trivial.

The conclusion of these reductions is that we have brought ourselves to the case where G is totally disconnected, and thus classical structure theory will not help us any further.

Let G^+ denote the discrete residual of G , i.e. the intersection of all open normal subgroups of G . By Theorem F in [2], G^+ is a characteristic cocompact subgroup of G and has no non-trivial discrete quotients. In particular, it is contained in every cocompact normal subgroup of G since all compact quotients of G are profinite. By Lemma 3, there exists an element in G/G^+ that normally generates it. Let gG^+ be the corresponding coset in G . Thus, no element of gG^+ is contained in any cocompact normal proper subgroup of G .

Every non-cocompact closed normal subgroup of G is contained in a maximal one, see Proposition 5.2 in [2]. It was proved in [2] that there are only finitely many such maximal non-cocompact closed normal subgroups in G — indeed, use Corollary 5.1 of [2] as in the proof of Theorem A ibidem. We denote them by M_1, \dots, M_k and assume that $k \geq 1$ since otherwise G is compact and we are done by Lemma 3. We claim that there are elements in the coset gG^+ which do not belong to any of the M_i . Such elements are therefore not contained in any proper closed normal subgroup of G ; that is, any such element normally generates G .

In order to prove the claim, suppose for a contradiction that gG^+ is contained in the union of all M_i . Then $gG^+ \cap M_i$ has non-empty interior in gG^+ for some i . Thus $G^+ \cap g^{-1}M_i$ has some interior point x relatively to G^+ . Therefore, the identity is an interior point of $G^+ \cap x^{-1}g^{-1}M_i$ within G^+ . Now $x^{-1}g^{-1}M_i = M_i$ and $G^+ \cap M_i$ is an open normal subgroup of G^+ . As G^+ has no non-trivial discrete quotients we conclude that $G^+ \cap M_i = G^+$ which means $G^+ < M_i$. This is absurd because G^+ is cocompact in G but M_i is not. This completes the proof of the claim and therefore of Theorem 1. \square

3. ON COMPACT GENERATION

Yves de Cornulier has kindly allowed us to present his example of a perfect locally compact group G which is not normally generated by a single element although it has no infinite discrete (group) quotient.

Assume that F is a finite perfect group with a subgroup $K < F$ such that

- (i) F is normally generated by K ,
- (ii) F is not normally generated by any single element of K .

Let G be the group of sequences in F that are ultimately in K . The “restricted product” locally compact group topology on G is defined by the identity neighbourhood base $U_n < K^{\mathbb{N}}$ consisting of sequences that are trivial on the first n coordinates. The perfect group $F^{(\mathbb{N})}$ of finitely supported sequences is then dense in G , so that G is perfect. Moreover, condition (ii) ensures that G cannot be normally generated by a single element.

Consider any discrete quotient of G . Its kernel must contain some U_n ; being normal, it follows by condition (i) that this quotient is a quotient of F^n , thus it is finite.

It remains to see that a pair $K < F$ as above indeed exists. Let c be an element of order two in a non-abelian finite simple group S . Set $F = S^3$ and let K be the (Klein) subgroup generated by $(1, c, c)$ and $(c, c, 1)$. Since the maximal normal subgroups of F are the kernels of the three canonical projections $F \rightarrow S$, which do not contain K , condition (i) holds. On the other hand, any element of K is in one of these kernels, whence condition (ii).

REFERENCES

- [1] G. Baumslag, A. G. Myasnikov, and V. Shpilrain. Open problems in combinatorial group theory. Second edition. In *Combinatorial and geometric group theory (New York, 2000/Hoboken, NJ, 2001)*, volume 296 of *Contemp. Math.*, pages 1–38. Amer. Math. Soc., Providence, RI, 2002.
- [2] P.-E. Caprace and N. Monod. Decomposing locally compact groups into simple pieces. *Math. Proc. Cambridge Philos. Soc.*, 150(1):97–128, 2011.
- [3] K. H. Hofmann and S. A. Morris. *The structure of compact groups*, volume 25 of *de Gruyter Studies in Mathematics*. Walter de Gruyter & Co., Berlin, augmented edition, 2006. A primer for the student—a handbook for the expert.
- [4] J. C. Lennox and J. Wiegold. Generators and killers for direct and free products. *Arch. Math. (Basel)*, 34(4):296–300, 1980.
- [5] V. D. Mazurov and E. I. Khukhro, editors. *The Kourovka notebook*. Russian Academy of Sciences Siberian Division Institute of Mathematics, Novosibirsk, augmented edition, 1999. Unsolved problems in group theory.
- [6] N. Monod, N. Ozawa, and A. Thom. Is an irng singly generated as an ideal? *Internat. J. Algebra Comput.*, 22(4):xx–xx, 2012.
- [7] D. Montgomery and L. Zippin. *Topological transformation groups*. Interscience Publishers, New York-London, 1955.

EPFL, SWITZERLAND